



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Short Communication

Long-term persistence of symptoms of dyspnoea in COVID-19 patients

Nhu Ngoc Nguyen^{1,2}, Van Thuan Hoang^{1,2,3}, Thi Loi Dao^{1,2,3}, Line Meddeb²,
Jean-Christophe Lagier^{2,4}, Matthieu Million^{2,4}, Didier Raoult^{2,4}, Philippe Gautret^{1,2,*}

¹ Aix Marseille University, IRD, AP-HM, SSA, VITROME, Marseille, France

² IHU-Méditerranée Infection, Marseille, France

³ Thai Binh University of Medicine and Pharmacy, Thai Binh, Vietnam

⁴ Aix Marseille University, IRD, AP-HM, MEPHI, Marseille, France

ARTICLE INFO

Article history:

Received 19 October 2021

Revised 19 November 2021

Accepted 22 November 2021

Keywords:

SARS-CoV-2

COVID-19

dyspnoea

long COVID

To the Editor

Even after recovering from the acute phase of COVID-19, patients may report the persistence of symptoms that have become known as 'long COVID' (Akbarialiabad *et al.*, 2021). In one meta-analysis of nine studies conducted on 1,816 patients between three weeks and three months after discharge from hospital, the persistence of dyspnoea, chest pain and a cough affected 37%, 16% and 14% of patients, respectively (Cares-Marambio *et al.*, 2021). These proportions gradually decrease with follow-up time. In a cross-sectional study conducted on 574 patients eight months after recovery, dyspnoea was the most frequent sequelae affecting 29% of patients (Zheng *et al.*, 2021). In one prospective cohort study conducted on 588 COVID-19 patients, 14% of patients reported dyspnoea at eight months of follow-up (Soraas *et al.*, 2021). In this study, we aim to estimate the rate of persistent dyspnoea in French COVID-19 patients, as evaluated after at least six months of follow-up and to investigate the risk factors for this persistence.

We identified confirmed COVID-19 patients who reported dyspnoea during the acute phase of the disease from among a cohort of 3,737 patients tested at our institute between 3 March and 27 April 2020 (Lagier *et al.*, 2020). Patients with pre-existing chronic respiratory diseases and chronic heart disease were excluded from this

analysis. Information on demographics, comorbidities, and acute symptoms were retrieved from medical files.

The selected patients were interviewed by telephone and asked to complete a questionnaire on possible persistent dyspnoea. Statistical analysis was performed by R 3.6.1 software (R Core team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, 2020. URL: <http://www.r-project.org>). Multiple logistic regression model were applied to explore the association between patient characteristics and dyspnoea at the acute phase and at the follow-up time. Variables with *p*-values < 0.2 in the univariable analysis were included in the multivariable analysis. Of the 3,737 COVID-19 patients, 3,222 had no known chronic respiratory disease and/or chronic heart disease at the time of inclusion and 880/3,222 (27.3%) reported dyspnoea at the acute phase of COVID-19. Being female or obese and consulting six or more days after the onset of symptoms were independently associated with dyspnoea. Patients with dyspnoea at admission were more likely to receive oxygen therapy. In univariate analysis, hypoxaemia, oxygen saturation $\leq 94\%$, C-reactive protein > 5 mg/L, neutrophils > 7.5 Giga/L, eosinophils < 0.1 Giga/L, lymphocyte < 1 Giga/L associated with dyspnoea, but a multivariate analysis was not conducted due to a high proportion of missing information. (Table 1). Of 880 patients, we randomly selected 838 patients for the telephone interview. Among them, 496/838 (59.2%) answered the questionnaire and 342/838 (40.8%) were lost to follow-up. Patients who responded were significantly less likely to report hypertension, to present severe symptoms, to require oxygen upon inclusion, to undergo prolonged hospitalisation and

* Corresponding author: Philippe Gautret, VITROME, Institut Hospitalo-Universitaire Méditerranée Infection, 19-21 Boulevard Jean Moulin 13385 Marseille Cedex 05, France.

E-mail address: philippe.gautret@club-internet.fr (P. Gautret).

Table 1
Risk factors for dyspnoea during the acute phase (n=3,222)

| | | No dyspnoea (n=2,342)% | Dyspnoea (n=880)% | Univariate analysis OR [95%CI] p-value | Multivariate analysis** OR [95%CI] p-value |
|--|---------------------|---------------------------|----------------------|---|---|
| Age | Mean \pm SD | 43.6 \pm 15.8 | 43.7 \pm 15.5 | | |
| | Range | 18–96 | 18–94 | | |
| | < 45 (n=1,720) | 1,255 (53.6) | 465 (52.8) | ref | |
| | \geq 45 (n=1,502) | 1,087 (46.4) | 415 (47.2) | 1.03 [0.88–1.21] 0.71 | |
| Sex | Male (n=1,464) | 1,107 (47.3) | 357 (40.6) | ref | ref |
| | Females (n=1,758) | 1,235 (52.7) | 523 (59.4) | 1.31 [1.12–1.54] 0.0007 | 1.38 [1.17–1.61] 0.0001 |
| Chronic conditions | | | | | |
| Hypertension | No (n=2,845) | 2,080 (88.8) | 765 (86.9) | ref | |
| | Yes (n=377) | 262 (11.2) | 115 (13.1) | 1.19 [0.94–1.52] 0.14 | |
| Diabetes | No (n=2999) | 2179 (93.0) | 820 (93.2) | ref | |
| | Yes (n=223) | 163 (7.0) | 60 (6.8) | 0.98 [0.71–1.34] 0.89 | |
| Cancer | No (n=3,138) | 2,280 (97.4) | 858 (97.5) | ref | |
| | Yes (n=84) | 62 (2.6) | 22 (2.5) | 0.94 [0.55–1.56] 0.81 | |
| Obesity | No (n=2,882) | 2,124 (90.7) | 758 (86.1) | ref | ref |
| | Yes (n=340) | 218 (9.3) | 122 (13.9) | 1.57 [1.23–1.99] 0.0002 | 1.45 [1.13–1.84] 0.003 |
| Co-medications | | | | | |
| Beta blockers | No (n=3,141) | 2,285 (97.6) | 856 (97.3) | ref | ref |
| | Yes (n=81) | 57 (2.4) | 24 (2.7) | 1.12 [0.66–1.85] 0.63 | |
| Dihydropyridine | No (n=3,117) | 2,269 (96.9) | 848 (96.4) | ref | |
| | Yes(n=105) | 73 (3.1) | 32 (3.6) | 1.17 [0.74–1.82] 0.46 | |
| Angiotensin-converting enzyme inhibitors | No (n=3,197) | 2,325 (99.3) | 872 (99.1) | ref | |
| | Yes (n=25) | 17 (0.7) | 8 (0.9) | 1.25 [0.47–3.08] 0.59 | |
| Angiotensin II receptor blocker | No (n=3,107) | 2,258 (96.4) | 849 (96.5) | ref | |
| | Yes (n=115) | 84 (3.6) | 31 (3.5) | 0.98 [0.62–1.51] 0.93 | |
| Metformin | No (n=3,125) | 2,273 (97.1) | 852 (96.8) | ref | |
| | Yes (n=97) | 69 (2.9) | 28 (3.2) | 1.08 [0.67–1.72] 0.72 | |
| Fenofibrate | No (n=3,203) | 2,329 (99.4) | 874 (99.3) | ref | |
| | Yes (n=19) | 13 (0.6) | 6 (0.7) | 1.23 [0.38–3.48] 0.68 | |
| Statin | No (n=3,154) | 2,290 (97.8) | 864 (98.2) | ref | |
| | Yes (n=68) | 52 (2.2) | 16 (1.8) | 0.82 [0.43–1.46] 0.48 | |

(continued on next page)

Table 1 (continued)

| | | No dyspnoea (n=2,342)% | Dyspnoea (n=880)% | Univariate analysis OR [95%CI] p-value | Multivariate analysis** OR [95%CI] p-value |
|--|------------------------|---------------------------|----------------------|---|---|
| COVID-19 status at inclusion | | | | | |
| Time between onset of COVID symptoms and admission | < 6 days(n=1,599) | 1,239 (52.9) | 360 (40.9) | ref | ref |
| | ≥ 6 days (n=1,623) | 1,103 (47.1) | 520 (52.1) | 1.62 [1.38–1.90] <0.0001 | 1.51 [1.28–1.77] <0.0001 |
| NEWS Score 2 | Low (n=3,017) | 2,216 (94.6) | 801 (91.0) | ref | |
| | Medium (n=125) | 89 (3.8) | 36 (4.1) | 1.12 [0.75–1.66] 0.58 | |
| | High (n=80) | 37 (1.6) | 43 (4.9) | 3.22 [2.06–5.03] <0.0001 | |
| Oxygen | No (n=3,034) | 2,256 (96.3) | 778 (88.4) | ref | |
| | Yes (n=188) | 86 (3.7) | 102 (11.6) | 3.44 [2.52–4.69] <0.0001 | 3.13 [2.31–4.25] <0.0001 |
| Oxygen saturation (SpO2) %* | >94% (n=2,754) | 1,968 (96.1) | 786 (92.4) | ref | |
| | ≤ 94% (n=145) | 80 (3.9) | 65 (7.6) | 2.03 [1.43–2.89] <0.0001 | |
| Chest Computed Tomography* | | | | | |
| | Normal (n=509) | 370 (32.2) | 139 (26.8) | ref | |
| | Limit (n=756) | 542 (47.1) | 214 (41.3) | 1.05 [0.82–1.35] 0.69 | |
| | Intermediate (n=329) | 203 (17.7) | 216 (24.3) | 1.65 [1.23–2.22] 0.001 | |
| | Severe (n=74) | 35 (3.0) | 39 (7.6) | 2.97 [1.81–4.87] <0.01 | |
| C-reactive protein * | ≤ 5mg/L (n=1,466) | 1,093 (63.2) | 373 (56.1) | ref | |
| | >5mg/L(n=929) | 637 (36.8) | 292 (43.9) | 1.34 [1.12–1.62] 0.001 | |
| Neutrophils* | ≤ 7.5 Giga/L (n=2,604) | 1,894 (97.8) | 710 (96.5) | ref | |
| | >7.5 Giga/L (n=68) | 42 (2.2) | 26 (3.5) | 1.65 [0.96–2.78] 0.046 | |
| Eosinophils* | ≥0.1 Giga/L (n=684) | 520 (26.9) | 164 (22.3) | ref | |
| | <0.1 Giga/L (n=1,988) | 1,416 (73.1) | 572 (77.7) | 1.28 [1.04–1.58] 0.015 | |
| Lymphocytes* | ≥ 1 Giga/L (n=2,370) | 1,731 (89.4) | 639 (86.8) | ref | |
| | < 1 Giga/L (n=302) | 205 (10.6) | 97 (13.2) | 1.28 [0.98–1.67] 0.06 | |
| D-Dimer* | ≤ 0.5 μg/ml (n=335) | 227 (58.7) | 108 (57.8) | ref | |
| | >0.5μg/ml (n=239) | 160 (41.3) | 79 (42.3) | 1.04 [0.72–1.50] 0.84 | |
| Fibrinogen* | ≤ 4g/L (n=290) | 216 (46.2) | 74 (35.2) | ref | |
| | >4 g/L (n=388) | 252 (53.8) | 136 (64.8) | 1.58 [1.11–2.24] 0.008 | |
| PCR Ct value < 16 at admission * | No (n=2,460) | 1,773 (93.3) | 687 (94.9) | ref | |
| | Yes (n=164) | 127 (6.7) | 37 (5.1) | 0.75 [0.50–1.11] 0.14 | |

NEWS Score 2: National Early Warning Score

*Oxygen saturation, Chest Computed Tomography, C-reactive protein, neutrophil, eosinophil, lymphocyte, D-Dimer, Fibrinogen and Ct <16 were not included in the multivariate due to >5% missing data

**only significant results are presented in the multivariate analysis

Table 2
Risk factors for dyspnoea at follow-up (n=469)

| | | No dyspnoea persistence (n=373)% | Dyspnoea persistence (n=123)% | Univariate analysis OR [95%CI] p-value | Multivariate analysis** OR [95%CI] p-value |
|--|-----------------------|----------------------------------|-------------------------------|--|--|
| Age | Mean \pm SD | 41.0 \pm 13.4 | 43.6 \pm 12.8 | | |
| | Range | 18–89 | 20–72 | | |
| | < 45 (n=285) | 226 (60.6) | 59 (48.0) | ref | |
| | \geq 45 (n=211) | 147 (39.4) | 64 (52) | 1.67 (1.08–2.57) | 1.74 [1.15–2.63] |
| Sex | Male (n=176) | 138 (37.0) | 38 (30.9) | ref | 0.009 |
| | Females (n=320) | 235 (63.0) | 85 (69.1) | 1.31 [0.83–2.09] | |
| | | | | 0.22 | |
| Chronic conditions | | | | | |
| Hypertension | No (n=451) | 343 (92.0) | 108 (87.8) | ref | |
| | Yes (n=45) | 30 (8.0) | 15 (12.2) | 1.59 [0.76–3.18] | |
| | | | | 0.16 | |
| Diabetes | No (n=471) | 353 (94.6) | 118 (95.9) | ref | |
| | Yes (n=25) | 20 (5.4) | 5 (4.1) | 0.75 [0.21–2.11] | |
| | | | | 0.57 | |
| Cancer | No (n=483) | 362 (97.1) | 121 (98.4) | ref | |
| | Yes (n=13) | 11 (2.9) | 2 (1.6) | 0.54 [0.06–2.55] | |
| | | | | 0.43 | |
| Obesity | No (n=431) | 327 (87.7) | 104 (84.6) | ref | |
| | Yes (n=65) | 46 (12.3) | 19 (15.5) | 1.29 [0.69–2.38] | |
| | | | | 0.37 | |
| Co-medications | | | | | |
| Beta blockers | No (n=486) | 367 (98.4) | 119 (96.8) | ref | ref |
| | Yes (n=10) | 6 (1.6) | 4 (3.3) | 2.05 [0.42–8.82] | |
| | | | | 0.26 | |
| Dihydropyridine | No (n=487) | 365 (97.9) | 122 (99.2) | ref | |
| | Yes(n=9) | 8 (2.1) | 1 (0.8) | 0.37 [0.008–2.84] | |
| | | | | 0.34 | |
| Angiotensin-converting enzyme inhibitors | No (n=493) | 372 (99.7) | 121 (98.4) | ref | |
| | Yes (n=3) | 1 (0.3) | 2 (1.6) | 6.15 [0.31–363.66] | |
| | | | | 0.09 | |
| Angiotensin II receptor blocker | No (n=484) | 364 (97.6) | 120 (97.6) | ref | |
| | Yes (n=12) | 9 (2.4) | 3 (2.4) | 1.01 [0.17–4.14] | |
| | | | | 0.99 | |
| Metformin | No (n=487) | 366 (98.1) | 121 (98.4) | ref | |
| | Yes (n=9) | 7 (1.9) | 2 (1.6) | 0.86 [0.09–4.62] | |
| | | | | 0.86 | |
| Fenofibrate | No (n=493) | 371 (99.5) | 122 (99.2) | ref | |
| | Yes (n=3) | 2 (0.5) | 1 (0.8) | 1.52 [0.03–29.42] | |
| | | | | 0.73 | |
| COVID-19 status at inclusion | | | | | |
| Time between onset of COVID symptoms and admission | < 6 days(n=207) | 146 (39.1) | 61 (49.6) | ref | ref |
| | \geq 6 days (n=289) | 227 (60.9) | 62 (50.4) | 0.65 [0.42–1.01] | 0.65 [0.43–0.99] |
| | | | | 0.04 | 0.045 |

(continued on next page)

Table 2 (continued)

| | | No dyspnoea persistence (n=373)% | Dyspnoea persistence (n=123)% | Univariate analysis OR [95%CI] p-value | Multivariate analysis** OR [95%CI] p-value |
|-------------------------------------|-------------------------|-------------------------------------|----------------------------------|---|---|
| NEWS Score 2 | Low (n=473) | 357 (95.7) | 116 (94.3) | ref | |
| | Medium (n=13) | 9 (2.4) | 4 (3.3) | 1.37 [0.41–4.52] 0.61 | |
| | High (n=10) | 7 (1.9) | 3 (2.4) | 1.32 [0.33–5.18] 0.69 | |
| Oxygen | No (n=467) | 353 (94.6) | 114 (92.7) | ref | |
| | Yes (n=29) | 20 (5.4) | 9 (7.3) | 1.39 [0.54–3.31] 0.42 | |
| Saturation oxygen (SpO2) % | > 94% (n=453) | 335 (93.3) | 118 (97.5) | ref | |
| | ≤94% (n=27) | 24 (6.7) | 3 (2.5) | 0.35 [0.07–1.20] 0.08 | |
| Chest Computed Tomography* | Normal (n=74) | 55 (29.9) | 19 (27.9) | ref | |
| | Limit (n=105) | 77 (41.9) | 28 (41.2) | 1.05 [0.53–2.07] 0.88 | |
| | Intermediate (n=58) | 40 (21.7) | 18 (26.5) | 1.30 [0.61–2.79] 0.50 | |
| | Severe (n=15) | 12 (6.5) | 3 (4.4) | 0.72 [0.18–2.84] 0.64 | |
| C-reactive protein* | ≤ 5mg/L (n=219) | 162 (60.0) | 57 (62.6) | ref | |
| | >5mg/L (n=142) | 108 (40.0) | 34 (37.4) | 0.89 [0.53–1.50] 0.66 | |
| Neutrophil* | ≤ 7.5 Giga/L (n=407) | 300 (97.1) | 107 (100.0) | ref | |
| | >7.5 Giga/L (n=9) | 9 (2.9) | 0 (0.0) | 0.00 [0.00–1.21] 0.07 | |
| Eosinophil* | ≥0.1 Giga/L (n=96) | 82 (26.5) | 14 (13.1) | ref | |
| | <0.1 Giga/L (n=320) | 227 (73.5) | 93 (86.9) | 2.40 [1.27–4.81] 0.004 | |
| Lymphocyte* | ≥ 1 Giga/L (n=376) | 281 (90.0) | 95 (88.8) | ref | |
| | < 1 Giga/L (n=40) | 28 (9.1) | 12 (11.2) | 1.27 [0.56–1.69] 0.51 | |
| D-Dimer* | ≤ 0.5 µg/ml (n=71) | 51 (73.9) | 20 (60.6) | ref | |
| | >0.5µg/ml (n=31) | 18 (26.1) | 13 (39.4) | 1.84 [0.69–4.84] 0.17 | |
| Fibrinogen* | ≤ 4g/L (n=36) | 28 (40.6) | 8 (30.8) | ref | |
| | >4 g/L (n=59) | 41 (59.4) | 18 (69.2) | 1.54 [0.54–4.66] 0.38 | |
| PCR Ct value < 16 at admission * | No (n=396) | 297 (93.7) | 99 (95.2) | ref | |
| | Yes (n=25) | 20 (6.3) | 5 (4.8) | 0.75 [0.21–2.13] 0.57 | |
| Viral shedding ≥ 10 days* | No (n=285) | 220 (89.8) | 65 (83.3) | ref | |
| | Yes (n=38) | 25 (10.2) | 13 (11.7) | 1.76 [0.78–3.81] 0.12 | |
| Hospitalisation ≥ 10 days | No (n=485) | 366 (98.1) | 119 (97.6) | ref | |
| | Yes (n=10) | 7 (1.9) | 3 (2.5) | 1.32 [0.22–5.88] 0.69 | |

(continued on next page)

Table 2 (continued)

| | | No dyspnoea persistence (n=373)% | Dyspnoea persistence (n=123)% | Univariate analysis OR [95%CI] p-value | Multivariate analysis** OR [95%CI] p-value |
|--|--------------|----------------------------------|-------------------------------|--|--|
| Transfer to the ICU | No (n=485) | 364 (97.6) | 121 (98.4) | ref | |
| | Yes (n=11) | 9 (2.4) | 2 (1.6) | 0.69 [0.07–3.29] | 0.61 |
| Hydroxychloroquine + azithromycin ≥ 3 days | No (n= 46) | 35 (9.4) | 11 (8.9) | ref | |
| | Yes (n= 450) | 338 (90.6) | 112 (91.1) | 1.05 [0.50–2.38] | 0.88 |

NEW Score 2: National Early Warning Score

*Chest Computed Tomography, C-reactive protein, neutrophil, eosinophil, lymphocytes, D-Dimer, Fibrinogen, Ct < 16 and viral shedding were not included in the multivariate due to >5% missing data

**only significant results are presented in the multivariate analysis

had lower rates of mortality, as compared with all patients reporting dyspnoea at inclusion (Supplementary Table 1)

Among these 496 patients, 123 (24.8%) reported persistent dyspnoea. The duration between the onset of symptoms and the interview was 44.0 ± 10.3 weeks, ranging from 31.6 to 63.1 weeks. Being aged ≥ 45 years old and consulting early during the acute phase (less than six days after the onset of symptoms) were independently associated with persistent dyspnoea (Table 2).

Lung damage in COVID-19 patients could be explained by various underlying mechanisms, including viral and immune-mediated implications (Liu et al., 2020), which could cause the persistence of dyspnoea. Dyspnoea is, by nature, a highly subjective experience. Women are more prone to suffer dyspnoea, possibly due to a decreased surface area for pulmonary gas exchange relative to lung size-matched men (Cory et al., 2015). People who are obese have a decreased functional residual capacity and expiration reserve volume (Parameswaran et al., 2006). Therefore, it is not surprising that female and obese COVID-19 patients had a higher risk for dyspnoea at admission. We observed that dyspnoea was more frequent in patients consulting more than six days after the onset of symptoms, suggesting that it takes a few days for the virus to provoke such symptoms. The association with a high viral load upon admission suggests a direct cytopathic effect of viruses on lung tissue.

The association of symptom persistence with being older may suggest a lower capacity of cell regeneration due to ageing. We have no hypothesis, however, to explain why early consultation was associated with the persistence of dyspnoea.

The use of the telephone interview has some limitations, including the lack of respiratory functional tests to quantify dyspnoea levels and a lack of information about smoking status. In addition, patients lost to follow-up could potentially include a high proportion of patients who fully recovered their respiratory function. Despite these limitations, our study underlined the high prevalence of the persistence of dyspnoea (24.8%) in COVID-19 patients and its association with older age. Further investigation with a clinical evaluation of respiratory function over time is required in these patients.

Ethical approval

This study was approved by the Comité de Protection des Personnes Nord Ouest II (No. 2021-A01183-33) on 22/07/2021.

Consent for publication

All authors gave their consent for publication.

Availability of data and materials

All the data for this study will be made available upon reasonable request to the corresponding author.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

None declared.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Author contributions

Philippe Gautret and Didier Raoult devised the project, the main conceptual ideas. Nhu Ngoc Nguyen, Jean Christophe Lagier, Matthieu Million, Thi Loi Dao, Line Meddeb collected the data. Van Tuan Hoang and Nhu Ngoc Nguyen analyzed and interpreted data. Nhu Ngoc Nguyen and Philippe Gautret wrote the manuscript. All authors reviewed and approved the final version of the manuscript.

Philippe Gautret is the guarantor for this work.

Acknowledgements

Our thanks go to Yolande Obadia and all the doctors who volunteered to recruit patients by telephone interview at the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, including Pierre Bares, Robert Barrus, Souad Benali, Marc Benoit, Dominique Blanc, Christine Carissimi, Denis Charpin, Jean Delmont, Patricia Enel, Charbelle Ezzedine, Pascale Fournier, Marguerite François, Katell Guillon, Guy Marmottant, Marie-Christine Manca-Pelissier, Isabelle Pinot, Nicole Quintana Benyahya, Robert Loi, Dominique Ricci Cagnol, Nicole Sarles-Philip, Catherine Tamalet, Jean-Michel Vassault, Marie-Pierre Vergobbi, and Nicole Veschi.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.ijid.2021.11.035](https://doi.org/10.1016/j.ijid.2021.11.035).

References

- Akbarialiabad H, Taghrir MH, Abdollahi A, Ghahramani N, Kumar M, Paydar S, et al. Long COVID: a comprehensive systematic scoping review. *Infection* 2021.
- Cares-Marambio K, Montenegro-Jimenez Y, Torres-Castro R, Vera-Urbe R, Torralba Y, Alsina-Restoy X, et al. Prevalence of potential respiratory symptoms in survivors of hospital admission after coronavirus disease 2019 (COVID-19): a systematic review and meta-analysis. *Chron Respir Dis* 2021;18 14799731211002240.
- Cory JM, Schaeffer MR, Wilkie SS, Ramsook AH, Puyat JH, Arbour B, et al. Sex differences in the intensity and qualitative dimensions of exertional dyspnea in physically active young adults. *J Appl Physiol* (1985) 2015;119:998–1006.
- Lagier JC, Million M, Gautret P, Colson P, Cortaredona S, Giraud-Gatineau A, et al. Outcomes of 3,737 COVID-19 patients treated with hydroxychloroquine/azithromycin and other regimens in Marseille, France: a retrospective analysis. *Travel Med Infect Dis* 2020;36:101791.
- Liu J, Zheng X, Tong Q, Li W, Wang B, Sutter K, et al. Overlapping and discrete aspects of the pathology and pathogenesis of the emerging human pathogenic coronaviruses SARS-CoV, MERS-CoV, and 2019-nCoV. *J Med Virol* 2020;92:491–4.
- Parameswaran K, Todd DC, Soth M. Altered respiratory physiology in obesity. *Can Respir J* 2006.
- Soraas A, Bo R, Kalleberg KT, Ellingjord-Dale M, Landrø NI . Self-reported memory problems eight months after non-hospitalized COVID-19 in a large Cohort. *medRxiv* 2021.
- Zheng D, Zhang L, Feng B, Liu Y, Liu H, Hu J, et al. Long-term clinical sequelae and immunological features of COVID-19 survivors: a cross-sectional study in Wuhan, China. *Lancet* (preprint) 2021.